

Bileptons - Present Constraints and Future Prospects at Colliders

Martti Raidal¹

Department of Theoretical Physics, University of Valencia,
46100 Burjassot, Valencia, Spain

Abstract. Bosons which couple to two leptons and carry lepton number $L = 2$, generically called bileptons, appear in many extensions of the SM. We review the present constraints on bilepton parameters coming from low and high energy data. In particular, we point out that the amplitude of $\mu - e$ conversion in nuclei is enhanced by large logarithms if compared with the $\mu \rightarrow e\gamma$ one. Bilepton couplings will be tested most stringently at future colliders. For wide range of neutrino masses the cross section of one of the processes $e^-e^-(\mu^-\mu^-) \rightarrow \ell_i^-\ell_i^-$, $i = e, \mu, \tau$, mediated by scalar triplet bilepton, has such a *lower* bound that it should be observed at future facilities.

1. Introduction

One of the peculiar features of the standard model (SM) is that none of its bosons carry global quantum numbers. This is no longer the case in wide range of extensions of the SM, such as grand unified theories [1], theories with enlarged Higgs sectors [2], theories which generate neutrino Majorana masses [3] as well as technicolour theories [4] and theories of compositeness [5]. Following Ref. [6] bileptons are defined to be bosons which couple to two leptons (but not to quarks) and which carry two units of lepton number. They can be both scalars and vectors. Their interactions

¹ E-mail: raidal@titan.ific.uv.es

Table 1. Upper bounds on diagonal bilepton couplings for $m_B = 1$ TeV.

Process	Combination of couplings	Upper bound
Møller	λ^{ee}	4 (2)
$(g-2)_\mu$	$\lambda^{\mu\mu}$	10 (5)
$M - \bar{M}$	$\lambda^{ee}\lambda^{\mu\mu}$	0.2 (0.2)

need not necessarily conserve lepton flavour, but otherwise the symmetries of the SM should be respected. The most general $SU(2)_L \times U(1)_Y$ invariant renormalizable dimension four Lagrangian for bileptons is given by [6]

$$\begin{aligned}
\mathcal{L} = & -\lambda_1^{ij} L_1^+ (\bar{\ell}_i^c P_L \nu_j - \bar{\ell}_j^c P_L \nu_i) \\
& + \tilde{\lambda}_1^{ij} \tilde{L}_1^{++} \bar{\ell}_i^c P_R \ell_j \\
& + \lambda_2^{ij} L_{2\mu}^+ \bar{\nu}_i^c \gamma^\mu P_R \ell_j \\
& + \lambda_2^{ij} L_{2\mu}^{++} \bar{\ell}_i^c \gamma^\mu P_R \ell_j \\
& + \lambda_3^{ij} L_3^0 \bar{\nu}_i^c P_L \nu_j \\
& - \lambda_3^{ij} / \sqrt{2} L_3^+ (\bar{\ell}_i^c P_L \nu_j + \bar{\ell}_j^c P_L \nu_i) \\
& - \lambda_3^{ij} L_3^{++} \bar{\ell}_i^c P_L \ell_j \\
& + \text{h.c.} \quad ,
\end{aligned} \tag{1}$$

where the subscripts 1–3 label the dimension of the $SU(2)_L$ representation which the bileptons belong to and the indices $i, j = e, \mu, \tau$ stand for the lepton flavours. In this talk I assume the couplings λ^{ij} to be real. In that case λ is anti-symmetric for scalar singlets, symmetric for scalar triplets and arbitrary for vector bileptons.

In the following I shall concentrate on the doubly charged components of bilepton multiplets. First I shall review the present most stringent bounds on their parameters and later discuss their production at future colliders.

2. Present constraints

Due to large bilepton masses, $m_B \gtrsim \mathcal{O}(\infty)$ GeV (LEP II pair production threshold), and unknown couplings to leptons one can presently only constrain bilepton effective couplings of a generic form $G = \sqrt{2}\lambda^2/(8m_B^2)$. Negative results in searches for the lepton flavour violating processes $\ell_l \rightarrow 3\ell_f$ and $\ell_l \rightarrow \gamma\ell_f$, where $l = \mu, \tau$, and $f = e, \mu$, put orders of magnitude

Table 2. Upper bounds on bilepton couplings for $m_B = 1$ TeV from various low energy leptonic processes.

Process	Combination of couplings	Upper bound
$\mu \rightarrow 3e$	$\lambda^{\mu e} \lambda^{ee}$	$4(2) \cdot 10^{-5}$
$\mu \rightarrow e\gamma$	$(\lambda\lambda)^{\mu e}$	$1.2(1.2) \cdot 10^{-2}$
$\mu - e$	$(\lambda\lambda)^{\mu e}$	$7(7) \cdot 10^{-3}$

more stringent bounds on off-diagonal bilepton couplings than one obtains from Møller scattering, $(g - 2)_\mu$ studies as well as from the searches for muonium-antimuonium conversion for diagonal couplings λ^{ee} and $\lambda^{\mu\mu}$. To date there is no constraints on $\lambda^{\tau\tau}$ without involving off-diagonal elements.

Most stringent constraints on diagonal bilepton couplings [6, 7, 8] are summarized in Table 1. The limits are approximately equal for singlet and triplet scalars and given as the first numbers in the table while the limits for vector bileptons are presented in brackets. Here and in the following bilepton masses are always taken in units of 1 TeV. As can be seen, for the scale of new physics $\mathcal{O}(\infty)$ TeV only the muonium-antimuonium conversion experiment [7] constrains diagonal bilepton couplings in a meaningful way.

Most stringent constraints on combinations of couplings λ from other lepton flavour violating processes [6, 8, 9] are summarized in Table 2. For shortness we have presented the processes involving first two generations only. Constraints from the processes involving third generation are considerably weaker. As expected, the strongest limit derives from $\mu \rightarrow 3e$ since it can occur at tree level. However, it constrains only a particular combination of couplings. On the other hand, $\mu \rightarrow e\gamma$ and $\mu - e$ conversion limits apply on a sum of coupling constant products since all possible intermediate states can occur at loop level and, therefore, provide complementary information.

One should note here that contrary to the claim in Ref.[10] $\mu - e$ conversion is more sensitive to bilepton interactions than $\mu \rightarrow e\gamma$. The general $\bar{e}\mu\gamma$ vertex can be parametrized in terms of form factors as follows

$$j^\rho = \bar{e} \left[\gamma_\mu (f_{E0} + f_{M0}\gamma_5) \left(g^{\mu\rho} - \frac{q^\mu q^\rho}{q^2} \right) + (f_{M1} + f_{E1}\gamma_5) i\sigma^{\rho\nu} \frac{q_\nu}{m_\mu} \right] \mu, \quad (2)$$

where q is the momentum transferred by photon. While $\mu \rightarrow e\gamma$ is induced only by form factors proportional to $\sigma^{\rho\nu}$ term then $\mu - e$ conversion rate is proportional to $(|f_{E0} + f_{M1}|^2 + |f_{M0} + f_{E1}|^2)$. Since the form factors f_{E0}

and f_{M0} are enhanced by large $\ln(m_\ell^2/m_B^2) \sim \mathcal{O}(\infty)$, where m_ℓ stands for the charged lepton mass running in the loop, if compared with f_{E1} and f_{M1} then $\mu - e$ conversion is enhanced while $\mu \rightarrow e\gamma$ is not. The logarithmic factors arise from the diagrams in which photon is attached to the charged lepton. Therefore, one expects logarithmic enhancement of $\mu - e$ conversion also in models with, e.g., leptoquarks or broken R -parity but not in models considered in Ref.[10].

3. Production at colliders

The most stringent constraints on bilepton couplings and masses will be obtained at future linear and muon colliders [11]. Bileptons with masses $m_B \lesssim \sqrt{s}/2$ can be pair produced in e^+e^- collisions almost model-independently due to their couplings to photon [12]. Single production of bileptons in e^+e^- and $e^-\gamma$ collisions allows one to probe masses up to $\sqrt{s}$ and test all the couplings λ^{ij} more sensitively than any of the present low energy experiments [13]. The most appropriate for studying bileptons are e^-e^- and $\mu^-\mu^-$ collision modes in which resonant s-channel production of bileptons via the process

$$e^-e^-(\mu^-\mu^-) \rightarrow \ell_i^-\ell_i^-, \quad (3)$$

$i = e, \mu, \tau$, is possible [14, 15]. For realistic machine and beam parameters sensitivity of $\lambda^{ij} \lesssim 5 \cdot 10^{-5}$ can be achieved [15]. Vector bileptons can be discriminated from the scalar ones by studying photon distribution in $e^-e^- \rightarrow L^{--}\gamma$.

Despite of this extraordinary sensitivity it could happen that due to small λ 's or high bilepton masses no positive signal will be detected at future colliders. However, this may not be the case if neutrinos are massive as predicted by most of the extensions of the SM and the process (3) is mediated by triplet scalar bilepton.

If the sum of light neutrino masses exceeds ~ 90 eV at least one of them has to be unstable. In order not to overclose the Universe the lifetime and mass of such an unstable neutrino ν_l must satisfy the requirement [16]

$$\tau_{\nu_l} \lesssim 8.2 \cdot 10^{31} \text{ MeV}^{-1} \left(\frac{100 \text{ keV}}{m_{\nu_l}} \right)^2. \quad (4)$$

The radiative decay modes $\nu_l \rightarrow \nu_f \gamma$, $\nu_f \gamma \gamma$ are highly suppressed [17] and cannot satisfy the constraint (4). The same is also true for Z' contribution to $\nu_l \rightarrow 3\nu_f$ decay [18]. The only possibility which lefts over is the decay $\nu_l \rightarrow 3\nu_f$ due to neutrino mixings induced by the neutral component of triplet bilepton.

Clearly, since L^0 and L^{--} belong to the same multiplet the reaction (3) can be related to decay $\nu_l \rightarrow 3\nu_f$. Since the latter decay rate is bounded from below by the constraint (4) and limits on neutrino mixings also the cross section of (3) has a lower limit. Let us consider numerically the most conservative case, $\nu_\tau \rightarrow 3\nu_e$. From the constraints presented above one obtains the following bound on $G^{e\tau}$ which induces the process $e^-e^- \rightarrow \tau^-\tau^-$ [19]

$$G^{e\tau} \gtrsim 2 \cdot 10^{-3} \frac{\lambda_{\tau\tau}}{\lambda_{ee} + \lambda_{\tau\tau}} \text{TeV}^{-2}, \quad (5)$$

where $G^{e\tau} = \sqrt{2}\lambda^{ee}\lambda^{\tau\tau}/(8m_B^2)$. On the other hand, studies of the process (3) at linear colliders give that far off resonance the minimal testable G^{ij} are $G^{ij}(\text{min}) = 1.4 \cdot 10^{-4}/s \text{TeV}^{-2}$, approximately the same for all relevant i, j . Therefore, the process $e^-e^- \rightarrow \tau^-\tau^-$ should be detected at the 1 TeV linear collider unless $\lambda_{\tau\tau}/\lambda_{ee} \lesssim 10^{-1}$. On the other hand, if this is the case then

$$G^{ee} \gtrsim 2 \cdot 10^{-3} \text{TeV}^{-2}, \quad (6)$$

and the excess of the electron pairs due to the s-channel bilepton exchange will be detected. Note that the positive signal should be seen if $\sqrt{s} \gtrsim 0.3 \text{TeV}$ which is below the planned initial energy of the linear collider.

Similar argumentation applies to all possible neutrino decays with even higher lower limits. Therefore, if one of the neutrino mass, indeed, exceeds 90 eV one should detect at least one of the processes (3) at future colliders.

4. Conclusions

Bileptons appear in many realistic extensions of the SM. The present constraints on diagonal bilepton couplings are rather weak compared to the bounds on off-diagonal couplings. Some of the latter ones, $(\lambda\lambda)^{e\mu}$, will be tested about two orders of magnitude more stringently in planned $\mu - e$ conversion experiments. This process is most sensitive to the couplings due to logarithmic enhancement of the conversion amplitude.

Future linear and muon colliders will be more sensitive to bilepton couplings than any of the present experiments. In particular, if triplet scalar particles exist and some neutrino masses exceed 90 eV then the doubly charged component of the triplet should be detected in e^-e^- or $\mu^-\mu^-$ collision modes of the colliders.

Acknowledgments

I would like to thank Organizers for creating pleasant atmosphere at the Ringberg castle. I have benefitted from discussions with P. Herzceg and especially with F. Cuyper and A. Santamaria whom I thank warmly.

References

- [1] P.H. Frampton, B.H. Lee, *Phys. Rev. Lett.* **64** (1990) 619; P.B. Pal, *Phys. Rev. D* **43** (1991) 236; P. Frampton *Phys. Rev. Lett.* **69** (1992) 2889; F. Pisano, V. Pleitez *Phys. Rev. D* **46** (1992) 410; P. Frampton, *Mod. Phys. Lett. A* **7**, (1992) 2017.
- [2] H. Georgi, M. Machacek, *Nucl. Phys. B* **262** (1985) 463; K.S. Babu, V.S. Mathur, *Phys. Lett. B* **196** (1987) 218; J.C. Pati, A. Salam, *Phys. Rev. D* **10** (1974) 275; R.N. Mohapatra, J.C. Pati, *Phys. Rev. D* **11** (1975) 566, *ibid.* 2558; G. Senjanovic, R.N. Mohapatra, *Phys. Rev. D* **12** (1975) 1502; R.N. Mohapatra, G. Senjanovic, *Phys. Rev. D* **23** (1981) 165.
- [3] T.P. Cheng, L. Li, *Phys. Rev. D* **22** (1980) 2860; G.B. Gelmini, M. Roncadelli, *Phys. Lett. B* **99** (1981) 411; A. Zee, *Phys. Lett. B* **93** (1980) 389.
- [4] E. Fahri, L. Susskind, *Phys. Rep.* **74** (1981) 277; E. Eichten et al., *Rev. Mod. Phys.* **56** (1984) 579.
- [5] W. Buchmüller, *Acta Phys. Austr. Suppl XXVII* (1985) 517; B. Schrempp, Proc. of the XXIII Int. Conf. on High Energy Physics Berkeley.
- [6] F. Cuyppers, S. Davidson, MPI-PhT/96-45, PSI-PR-96-21 [hep-ph/9609487].
- [7] K. Jungman, these proceedings.
- [8] M.D. Swartz, *Phys. Rev. D* **40** (1989) 1521; J.F. Gunion et al., *Phys. Rev. D* **40** (1989) 1546; R. Mohapatra, *Phys. Rev. D* **46** (1992) 2990.
- [9] M. Raidal and A. Santamaria, FTUV/97-56, to be published.
- [10] See, e.g., T.S. Kosmas and J.D. Vergados, *Phys. Rev.* **164** (1996) 251; and references therein.
- [11] J.F. Gunion, UCD-97-17 [hep-ph/9707379].
- [12] J.F. Gunion, C. Loomis and K.T. Pitts, UCD-96-30 [hep-ph/9610237].
- [13] N. Lepore, B. Thorndyke, H. Nadeau, D. London, *Phys. Rev. D* **50** (1994) 2031; G. Barenboim, K. Huitu, J. Maalampi, and M. Raidal, *Phys. Lett. B* **394** (1997) 132; and references therein.
- [14] T. Rizzo, *Phys. Rev. D* **25** (1982) 1355; T. Rizzo, *Phys. Rev. D* **46** (1992) 910; P. Frampton, D. Ng, *Phys. Rev. D* **45** (1992) 4240; J. Maalampi, A. Pietilä, M. Raidal, *Phys. Rev. D* **48** (1993) 4467.
- [15] J.F. Gunion, *Int. J. Mod. Phys. A* **11** (1996) 1551; F. Cuyppers and M. Raidal, PSI-PR-97-11, FTUV/97-13 [hep-ph/9704224].
- [16] E. Kolb and M. Turner, *Phys. Rev. Lett.* **67** (1991) 5; P. Herczeg and R.N. Mohapatra, *Phys. Rev. Lett.* **69** (1992) 2475.
- [17] A. De Rujula and S.L. Glashow, *Phys. Rev. Lett.* **45** (1980) 942; P.B. Pal and L. Wolfenstein, *Phys. Rev. D* **25** (1982) 766.
- [18] Y. Hosotani, *Nucl. Phys. B* **191** (1981) 411; J. Schechter and J.W. Valle, *Phys. Rev. D* **25** (1982) 774.
- [19] M. Raidal, FTUV/97-34, [hep-ph/9706279].